

# Suppression of convective precipitation by elevated man-made aerosols is responsible for large-scale droughts in north China

Zhen Li<sup>a,b</sup>, Shaocai Yu<sup>a,b,c,1</sup>, Liqiang Wang<sup>a,b</sup>, Khalid Mehmood<sup>a,b</sup>, Weiping Liu<sup>a,b</sup>, and Kiran Alapaty<sup>d</sup>

It has been proposed that the summer “South Flood–North Drought” (SFND) pattern observed in China over recent decades is caused by the relative impacts of global warming, aerosol loading, and natural variability on regional rainfall (1–3). This conclusion is supported by a recent study by Day et al. (4) in which the SFND is attributed to the changes in the frequency of frontal rain events. Using a technique called the Frontal Rain Event Detection Algorithm for the observations during 1951–2007, decadal changes in the amount and distribution of rainfall in eastern China were found to be overwhelmingly due to changes in frontal rainfall (4). Day et al. conclude that frontal rainfall was envisioned as the product of large-scale frontal convergence and the nonfrontal rainfall was because of local convection, orographic rainfall, and typhoon rainfall. The authors further imply that the shifts in frequency and latitude of frontal rainfall over the recent decades in eastern China reflect changes in large-scale atmospheric circulation.

However, Day et al.’s (4) conclusion fails to consider suppression of convective precipitation in north China by anthropogenic aerosols, a critically important factor for the recent SFND in China (5–7). On the basis of the observed summer mean rainfall over China during 1952–2015, Yu et al. (7) found that large-scale rainfall enhancements are persistently occurring in east China (along 27.0°N to 34.5°N, mainly located in Yangtze River Delta) with increasing droughts in north China (along the 34.5°N to 45°N latitudinal band). By cooling the surface and the lower troposphere over the land and stabilizing the boundary layer, the enhanced

anthropogenic aerosols suppressed the deep convection, which reduces the condensational heating in the free troposphere and the thermal contrast between the land and ocean over north China (5, 6). The enhancement of the large-scale precipitation in east China and the suppression of the convective precipitation in north China by the elevated anthropogenic aerosols are believed to be responsible for the SFND (5–7).

Although we strongly agree with Day et al.’s (4) finding that decadal changes in rainfall in eastern China (mainly located in Yangtze River Delta) are mainly due to changes in frontal rainfall because of the enhancement of the large-scale precipitation, we do not believe that the large-scale droughts persistently occurring in north China (along the 34.5°N to 45°N latitudinal band) are mainly due to a decrease of frontal rain event frequency. We believe that the suppression of the convective precipitation (i.e., nonfrontal rain) in north China by the elevated anthropogenic aerosols is responsible for this. Further research with the observations and modeling effort is urgent.

## Acknowledgments

This work was partially supported by Department of Science and Technology of China Grants 2016YFC0202702 and 2014BAC22B06 and National Natural Science Foundation of China Grant 21577126. This work was also supported by the Joint National Natural Science Foundation of China–Israel Science Foundation Research Program Grant 41561144004. Part of this work was also supported by the “Zhejiang 1000 Talent Plan” and Research Center for Air Pollution and Health in Zhejiang University. The views expressed in this paper are those of the author(s) and do not necessarily represent those of the US Environmental Protection Agency.

- 1 Broecker WS, Putnam AE (2013) Hydrologic impacts of past shifts of Earth’s thermal equator offer insight into those to be produced by fossil fuel CO<sub>2</sub>. *Proc Natl Acad Sci USA* 110:16710–16715.
- 2 Bollasina MA, Ming Y, Ramaswamy V (2011) Anthropogenic aerosols and the weakening of the South Asian summer monsoon. *Science* 334:502–505.

<sup>a</sup>Key Laboratory of Environmental Remediation and Ecological Health, Ministry of Education, College of Environmental and Resource Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, People’s Republic of China; <sup>b</sup>Research Center for Air Pollution and Health, College of Environmental and Resource Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, People’s Republic of China; <sup>c</sup>Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA 91125; and <sup>d</sup>Systems Exposure Division, National Exposure Research Laboratory, US Environmental Protection Agency, Research Triangle Park, NC 27711

Author contributions: S.Y. designed research; Z.L., S.Y., L.W., K.M., W.L., and K.A. performed research; Z.L., S.Y., L.W., K.M., W.L., and K.A. analyzed data; and Z.L. and S.Y. wrote the paper.

The authors declare no conflict of interest.

Published under the [PNAS license](#).

<sup>1</sup>To whom correspondence should be addressed. Email: shaocaiyu@zju.edu.cn.

Published online August 21, 2018.

- 3 Menon S, Hansen J, Nazarenko L, Luo Y (2002) Climate effects of black carbon aerosols in China and India. *Science* 297:2250–2253.
- 4 Day JA, Fung I, Liu W (2018) Changing character of rainfall in eastern China, 1951–2007. *Proc Natl Acad Sci USA* 115:2016–2021.
- 5 Jiang YQ, Liu XH, Yang XQ, Wang MH (2013) A numerical study of the effect of different aerosol types on East Asian summer clouds and precipitation. *Atmos Environ* 70:51–63.
- 6 Wu GX, et al. (2016) Advances in studying interactions between aerosols and monsoon in China. *Sci China Earth Sci* 59:1–16.
- 7 Yu S, et al. (2016) Anthropogenic aerosols are a potential cause for migration of the summer monsoon rain belt in China. *Proc Natl Acad Sci USA* 113:E2209–E2210.